arXiv:1602.08659v1 [math.NT] 28 Feb 2016

DIFFERENTIAL EQUATIONS FOR CHANGHEE POLYNOMIALS AND THEIR APPLICATIONS

DMITRY V. DOLGY, DAE SAN KIM, TAEKYUN KIM, JONG JIN SEO

Abstract. Recently, the non-linear Changhee differential equations were introduced in [5] and these differential equations turned out to be very useful for studying special polynomials and mathematical physics. Some interesting identities and properties of Changhee polynomials can also be derived from umbral calculus (see [7]). In this paper, we consider differential equations arising from Changhee polynomials and derive some new and explicit formulae and identities from our differential equations.

1. Introduction

As is well known, the Euler polynomials are defined by the generating function

$$\frac{2}{e^t+1}e^{xt}=\sum_{n=0}^{\infty}E_n(x)\frac{t^n}{n!},\ (\text{see } [1,3,10]). \tag{1.1}$$

With the viewpoint of deformed Euler polynomials, the Changhee polynomials are defined by the generating function

$$\frac{2}{2+t}(1+t)^x=\sum_{n=0}^{\infty}Ch_n(x)\frac{t^n}{n!},\ (\text{see } [3]). \tag{1.2}$$

2010 *Mathematics Subject Classification.* 05A19, 11B83, 34A30.
Key words and phrases. Changhee polynomials, differential equations.

.

From (1.2), we note that

$$\begin{aligned}\frac{2}{e^{log(1+t)}+1}e^{xlog(1+t)} &= \sum_{n=0}^{\infty} E_n(x)\frac{1}{n!}(log(1+t))^n \\ &= \sum_{n=0}^{\infty} E_n(x)\sum_{m=n}^{\infty} S_1(m,n)\frac{t^m}{m!} \\ &= \sum_{m=0}^{\infty}\left(\sum_{n=0}^{m} E_n(x)S_1(m,n)\right)\frac{t^m}{m!},\end{aligned} \tag{1.3}$$

where $S_1(m,n)$ is the stirling number of the first kind which is defined as

$$(x)_0 = 1, (x)_n = x(x-1)\cdots(x-n+1) = \sum_{l=0}^{n} S_1(n,l)x^l, (n \geq 1). \tag{1.4}$$

From (1.2) and (1.3), we note that

$$Ch_m(x) = \sum_{n=0}^{m} E_n(x)S_1(m,n), (m \geq 0), (\text{see } [3,12,13]). \tag{1.5}$$

Replacing t by $e^t - 1$ in (1.2), we get

$$\begin{aligned}\sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!} = \frac{2}{e^t+1}e^{xt} &= \sum_{m=0}^{\infty} Ch_m(x)\frac{1}{m!}(e^t-1)^m \\ &= \sum_{m=0}^{\infty} Ch_m(x)\sum_{n=m}^{\infty} S_2(n,m)\frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty}\left(\sum_{m=0}^{n} Ch_m(x)S_2(n,m)\right)\frac{t^n}{n!},\end{aligned} \tag{1.6}$$

where $S_2(n,m)$ is the stirling number of the second kind which is given by $x^n = \sum_{l=0}^{n} S_2(n,l)(x)_l, (n \geq 0)$. Thus, by (1.6), we get

$$E_n(x) = \sum_{m=0}^{n} Ch_m(x)S_2(n,m), (n \geq 0), (\text{see } [3]). \tag{1.7}$$

Recently, several authors have studied Changhee polynomials and numbers (see [1-20]). In this paper, we consider differential equations derived from the generating function of Changhee polynomials and give some new and explicit formulae for the Changhee polynomials by using our results on differential equations.

2. Differential equations for Chagnhee polynomials

Let

$$F = F(t,x) = \frac{1}{2+t}(1+t)^x. \tag{2.1}$$

From (2.1), we can derive the following equations:

$$F^{(1)} = \frac{d}{dt}F(t,x) = (-(2+t)^{-1} + x(1+t)^{-1})F, \tag{2.2}$$

$$F^{(2)} = \frac{d}{dt}F^{(1)} = (2(2+t)^{-2} - 2x(2+t)^{-1}(1+t)^{-1} + (x^2-x)(1+t)^{-2})F, \tag{2.3}$$

and

$$\begin{aligned} F^{(3)} = \frac{d}{dt}F^{(2)} &= (-6(2+t)^{-3} + 6x(2+t)^{-2}(1+t)^{-1} \\ &\quad + (3x-3x^2)(2+t)^{-1}(1+t)^{-2} + (x^3-3x^2+2x)(1+t)^{-3})F. \end{aligned} \tag{2.4}$$

Continuing this process, we put

$$F^{(N)} = \left(\frac{d}{dt}\right)^N F(t,x) = (\sum_{i=0}^{N} a_i(N,x)(1+t)^{-i}(2+t)^{i-N})F, \tag{2.5}$$

where $N = 0, 1, 2, \cdots$. From (2.5), we note that

$$\begin{aligned} F^{(N+1)} = \frac{d}{dt}F^{(N)} &= \left(\sum_{i=0}^{N} a_i(N,x)(-i)(1+t)^{-i-1}(2+t)^{i-N}\right)F \\ &\quad + \left(\sum_{i=0}^{N} a_i(N,x)(1+t)^{-i}(i-N)(2+t)^{i-N-1}\right)F \\ &\quad + \left(\sum_{i=0}^{N} a_i(N,x)(1+t)^{-i}(2+t)^{i-N}\right)F^{(1)} \\ &= \Bigg\{\sum_{i=0}^{N}(-i)a_i(N,x)(1+t)^{-i-1}(2+t)^{i-N} \\ &\quad + \sum_{i=0}^{N}(i-N)a_i(N,x)(1+t)^{-i}(2+t)^{i-N-1} \\ &\quad - \sum_{i=0}^{N} a_i(N,x)(1+t)^{-i}(2+t)^{i-N-1} \\ &\quad + \sum_{i=0}^{N} x a_i(N,x)(1+t)^{-i-1}(2+t)^{i-N}\Bigg\}F \end{aligned} \tag{2.6}$$

$$= \Bigg\{ \sum_{i=1}^{N+1} (1-i) a_{i-1}(N,x)(1+t)^{-i}(2+t)^{i-N-1} + \sum_{i=0}^{N} (i-N) a_i(N,x)(1+t)^{-i}(2+t)^{i-N-1} - \sum_{i=0}^{N} a_i(N,x)(1+t)^{-i}(2+t)^{i-N-1} + \sum_{i=1}^{N+1} x a_{i-1}(N,x)(1+t)^{-i}(2+t)^{i-N-1} \Bigg\} F.$$

On the other hand, by replacing N by $N+1$ in (2.5), we get

$$F^{(N+1)} = \{ \sum_{i=0}^{N+1} a_i(N+1,x)(1+t)^{-i}(2+t)^{i-N-1} \} F. \tag{2.7}$$

Comparing the coefficients on both sides of (2.6) and (2.7), we have

$$a_0(N+1,x) = -(N+1) a_0(N,x), \tag{2.8}$$

$$a_{N+1}(N+1,x) = (x-N) a_N(N,x), \tag{2.9}$$

and

$$a_i(N+1,x) = (x+1-i) a_{i-1}(N,x) + (i-N-1) a_i(N,x), \tag{2.10}$$

where $1 \le i \le N$.
We also note that

$$F = F^{(0)} = a_0(0,x) F. \tag{2.11}$$

Thus, by (2.11), we get

$$a_0(0,x) = 1. \tag{2.12}$$

From (2.2) and (2.5), we can derive the following equation:

$$\begin{aligned} (-(2+t)^{-1} + x(1+t)^{-1}) F = F^{(1)} &= \left(\sum_{i=0}^{1} a_i(1,x)(1+t)^{-i}(2+t)^{i-1} \right) F \\ &= \left(a_0(1,x)(2+t)^{-1} + a_1(1,x)(1+t)^{-1} \right) F. \end{aligned} \tag{2.13}$$

Comparing the coefficients on both sides of (2.13), we get

$$a_0(1,x) = -1, a_1(1,x) = x. \tag{2.14}$$

Also, by (2.8) and (2.9), we have

$$\begin{aligned} a_0(N+1,x) &= -(N+1) a_0(N,x) = (-1)^2 (N+1) N a_0(N-1,x) = \cdots \\ &= (-1)^N (N+1) N \cdots 2 a_0(1,x) = (-1)^{N+1} (N+1)!, \end{aligned} \tag{2.15}$$

and

$$\begin{aligned} a_{N+1}(N+1,x) &= (x-N)a_N(N,x) = (x-N)(x-(N-1))a_{N-1}(N-1,x) \\ &= \cdots = (x-N)(x-(N-1))\cdots(x-1)a_1(1,x) \\ &= (x-N)(x-(N-1))\cdots(x-1)x = (x)_{N+1}. \end{aligned} \tag{2.16}$$

From (2.10), we can derive the following equations:

$$\begin{aligned} a_1(N+1,x) &= xa_0(N,x) - Na_1(N,x) \\ &= x\left(a_0(N,x) - Na_0(N-1,x)\right) + (-1)^2N(N-1)a_1(N-1,x) \\ &= \cdots \\ &= x\sum_{i=0}^{N-1}(-1)^i(N)_i a_0(N-i,x) + (-1)^N N!a_1(1,x) \\ &= x\sum_{i=0}^{N}(-1)^i(N)_i a_0(N-i,x), \end{aligned} \tag{2.17}$$

$$\begin{aligned} a_2(N+1,x) &= (x-1)a_1(N,x) + (1-N)a_2(N,x) \\ &= (x-1)\left(a_1(N,x) + (-1)(N-1)a_1(N-1,x)\right) \\ &\quad + (-1)^2(N-1)(N-2)a_2(N-1,x) \\ &= \cdots \\ &= (x-1)\sum_{i=0}^{N-2}(-1)^i(N-1)_i a_1(N-i,x) + (-1)^{N-1}(N-1)!a_2(2,x) \\ &= (x-1)\sum_{i=0}^{N-1}(-1)^i(N-1)_i a_1(N-i,x), \end{aligned} \tag{2.18}$$

and

$$\begin{aligned} a_3(N+1,x) &= (x-2)a_2(N,x) + (2-N)a_3(N,x) \\ &= (x-2)\left(a_2(N,x) + (-1)(N-2)a_2(N-1,x)\right) \\ &\quad + (-1)^2(N-2)(N-3)a_3(N-1,x) \\ &= \cdots \end{aligned} \tag{2.19}$$

$$= (x-2)\sum_{i=0}^{N-3}(-1)^i(N-2)_i a_2(N-i,x) + (-1)^{N-2}(N-2)!a_3(3,x)$$
$$= (x-2)\sum_{i=0}^{N-2}(-1)^i(N-2)_i a_2(N-i,x).$$

Continuing this process, we have

$$a_j(N+1,x) = (x-j+1)\sum_{i=0}^{N-j+1}(-1)^i(N-j+1)_i a_{j-1}(N-i,x), (1\le j\le N). \tag{2.20}$$

The matrix $(a_i(j,x))_{0\le i,j\le N}$ is given by

$$\begin{array}{c} \\ 0 \\ 1 \\ 2 \\ \vdots \\ N \end{array}\begin{array}{c} \begin{array}{cccccc} 0 & 1 & 2 & \dots & N \end{array} \\ \begin{pmatrix} 1 & -1! & (-1)^2 2! & \dots & (-1)^N N! \\ 0 & x & \dots & \dots & \cdot \\ 0 & 0 & (x)_2 & \dots & \cdot \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & (x)_n \end{pmatrix} \end{array}$$

Now, we give explicit expressions for $a_i(j,x)$. From (2.15), (2.17), (2.18), (2.19) and (2.20), we can derive the following equations:

$$a_1(N+1,x) = x\sum_{i=0}^{N}(-1)^i(N)_i a_0(N-i,x) = x(-1)^N(N+1)!, \tag{2.21}$$

$$\begin{aligned} a_2(N+1,x) &= (x-1)\sum_{i_1=0}^{N-1}(-1)^{i_1}(N-1)_{i_1}a_1(N-i_1,x) \\ &= (x)_2(-1)^{N-1}(N-1)!\sum_{i_1=0}^{N-1}(N-i_1), \end{aligned} \tag{2.22}$$

$$\begin{aligned} a_3(N+1,x) &= (x-2)\sum_{i_2=0}^{N-2}(-1)^{i_2}(N-2)_{i_2}a_2(N-i_2,x) \\ &= (x)_3(-1)^{N-2}(N-2)!\sum_{i_2=0}^{N-2}\sum_{i_1=0}^{N-2-i_2}(N-i_2-i_1-1), \end{aligned} \tag{2.23}$$

and

$$a_4(N+1,x) = (x-3)\sum_{i_3=0}^{N-3}(-1)^{i_3}(N-3)_{i_3}a_3(N-i_3,x)$$
$$= (x)_4(-1)^{N-3}(N-3)!\sum_{i_3=0}^{N-3}\sum_{i_2=0}^{N-3-i_3}\sum_{i_1=0}^{N-3-i_3-i_2}(N-i_3-i_2-i_1-2). \tag{2.24}$$

Continuing this process, we get

$$a_j(N+1,x) = (x)_j(-1)^{N-j+1}(N-j+1)!$$
$$\times\sum_{i_{j-1}=0}^{N-j+1}\sum_{i_{j-2}=0}^{N-j+1-i_{j-1}}\cdots\sum_{i_1=0}^{N-j+1-i_{j-1}-\cdots-i_2}(N-i_{j-1}\cdots-i_1-j+2), \tag{2.25}$$

where $1 \le j \le N+1$.
Therefore, by (2.25), we obtain the following theorem.

Theorem 2.1. *For $N = 0, 1, 2, \cdots$, the linear differential equations*

$$F^{(N)} = \left(\sum_{i=0}^{N} a_i(N,x)(1+t)^{-i}(2+t)^{i-N}\right)F$$

has a solution $F = F(t,x) = \frac{1}{2+t}(1+t)^x$,
where

$$a_0(N,x) = (-1)^N N!,$$
$$a_j(N,x) = (x)_j(-1)^{N-j}(N-j)!$$
$$\times\sum_{i_{j-1}=0}^{N-j}\sum_{i_{j-2}=0}^{N-j-i_{j-1}}\cdots\sum_{i_1=0}^{N-j-i_{j-1}-\cdots-i_2}(N-i_{j-1}\cdots-i_1-j+1), (1 \le j \le N).$$

We recall that Changhee polynomials, $Ch_n(x), (n \ge 0)$, are given by the generating function

$$2F = 2F(t,x) = \frac{2}{2+t}(1+t)^x = \sum_{n=0}^{\infty} Ch_n(x)\frac{t^n}{n!}. \tag{2.26}$$

On the one hand, by (2.26), we get

$$2F^{(N)} = \sum_{k=N}^{\infty} Ch_k(x)(k)_N\frac{t^{k-N}}{k!} = \sum_{k=0}^{\infty} Ch_{N+k}(x)\frac{t^k}{k!}. \tag{2.27}$$

On the other hand, by Theorem 1, we have

$$\begin{aligned}
2F^{(N)} &= \left(2\sum_{i=0}^{N} a_i(N,x)(1+t)^{-i}(2+t)^{i-N}\right)F \\
&= \sum_{i=0}^{N} 2a_i(N,x)\left(\sum_{l=0}^{\infty}(-1)^l\binom{i+l-1}{l}t^l\right) \\
&\quad\times\left(\sum_{m=0}^{\infty}(-1)^m 2^{i-N-m}\binom{N+m-i-1}{m}t^m\right)\left(\sum_{p=0}^{\infty}Ch_p(x)\frac{t^p}{p!}\right) \\
&= \sum_{i=0}^{N} a_i(N,x)\sum_{k=0}^{\infty}\sum_{l+m+p=k}(-1)^{l+m}2^{i-N-m+1}\binom{i+l-1}{l} \qquad (2.28)\\
&\qquad\times\binom{N+m-i-1}{m}\frac{1}{p!}Ch_p(x)t^k \\
&= \sum_{k=0}^{\infty}\left\{k!\sum_{i=0}^{N}a_i(N,x)\sum_{l+m+p=k}(-1)^{l+m}\frac{2^{i-N-m+1}}{p!}\binom{l+i-1}{l}\right. \\
&\qquad\left.\binom{N+m-i-1}{m}Ch_p(x)\right\}\frac{t^k}{k!}.
\end{aligned}$$

By comparing the coefficients on the both sides of (2.27) and (2.28), we obtain the following theorem

Theorem 2.2. *For $k, N = 0, 1, 2, \cdots$, we have*

$$\begin{aligned}
Ch_{k+N}(x) &= k!\sum_{i=0}^{N}a_i(N,x)\sum_{l+m+p=k}(-1)^{l+m}\frac{2^{i-N-m+1}}{p!}\binom{l+i-1}{l} \\
&\qquad\times\binom{N+m-i-1}{m}Ch_p(x) \\
&= \sum_{i=0}^{N}a_i(N,x)\sum_{l+m+p=k}(-1)^{l+m}2^{i-N-m+1} \\
&\qquad\times\binom{k}{l,m,p}(i+l-1)_l(N+m-i-1)_m Ch_p(x),
\end{aligned}$$

where

$$a_0(N,x) = (-1)^N N!,$$

$$a_j(N,x) = (x)_j(-1)^{N-j}(N-j)! \times \sum_{i_{j-1}=0}^{N-j} \sum_{i_{j-2}=0}^{N-j-i_{j-1}} \cdots \sum_{i_1=0}^{N-j-i_{j-1}-\cdots-i_2} (N-i_{j-1}-\cdots-i_1-j+1),$$

$$(1 \le j \le N).$$

References

1. A. Bayad, J. Chikhi, *Apostol-Euler polynomials and asymptotics for negative binomial reciprocals,* Adv. Stud. Contemp. Math. (Kyungshang) **24** (2014), no. 1, 33-37.
2. L. C. Jang, C. S. Ryoo. J. J. Seo, H. K. kwon, *Some properties of the twisted Changhee polynomials and their zeros,* Appl. Math. Comput. **274** (2016), no. 1, 169-177.
3. D. S. Kim, T. Kim, *A note on Boole polynomials,* Integral Transforms Spec. Funct. **25** (2014), no. 8, 627-633.
4. D. S. Kim, T. Kim, *Some identities of Korobov-type polynomials associated with p-adic integrals on $\mathbb{Z}_p$,* Adv. Difference Equ. **2015** 2015:282.
5. D. S. Kim, T. Kim, *A note on non-linear Changhee differential equations,* Russ. J. Math. Phys. (to appear).
6. D. S. Kim, T. Kim, T. Komatsu, S.-H. Lee, *Barnes-type Daehee of the first kind and poly-Cauchy of the first kind mixed-type polynomials,* Adv. Difference Equ. **2014** 2014:140.
7. D. S. Kim, T. Kim, J. J. Seo, *A Note on Changhee Polynomials and Numbers,* Adv. Studies Theor. Phys. **7** (2013), no. 20, 993-1003.
8. T. Kim, *Non-Archimedean q-integrals associated with multiple Changhee q-Bernoulli polynomials,* Russ. J. Math. Phys. **10** (2003), no. 1, 91-98.
9. T. Kim, T. Mansour, *Umbral calculus associated with Frobenius-type Eulerian polynomials* Russ. J. Math. Phys. **21** (2014), no. 4, 484-493.
10. D. V. Kruchinin, V. V. Kruchinin, *Application of a composition of generating functions for obtaining explicit formulas of polynomials,* J. Math. Anal. Appl. **404** (2013), no. 1, 161-171.
11. H. I. Kwon, T. Kim, J.-J. Seo, *A note on degenerate Changhee numbers and polynomials,* Proc. Jangjeon Math.Soc. **18** (2015), no. 1, 295-305.
12. J. Kwon, H. S. Noh, S. H. Jeong, A. J. Kim, J. H. Lee, S. H. Rim, *A Note on weighted Changhee Polynomials and Numbers,* Adv. Studies. Theor. Phys. **9** (2015), no. 4, 191-198.
13. D. Lim, F. Qi, *On the Appell type λ-Changhee polynomials,* J. Nonlinear Sci. Appl. **9** (2016), 1872-1876.
14. J.-W. Park, *On the twisted q-Changhee polynomials of higher order,* J. Comput. Anal. Appl. **20** (2016), no. 3, 424-431.
15. S.-H. Rim, J.-W. Park, S.-S. Pyo, J. Kwon, *The n-th twisted Changhee polynomials and numbers,* Bull. Korean Math. Soc. **52** (2015), no. 3, 741-749.
16. C. S. Ryoo, T. Kim, R. P. Agarwal, *Exploring the multiple Changhee q-Bernoulli polynomials,* Int. J. Comput. Math. **82** (2005), no. 4, 483-493.

17. C. S. Ryoo, H. Song, R. P. Agarwal, *On the roots of the q-analogue of Euler-Barnes' polynomials,* Adv. Stud. Contemp. Math. (Kyungshang) **9** (2004), no. 2, 153-163.
18. G. Y. Sohn, J. K. Kwon, *A note on twisted Changhee polynomials and numbers with weight,* Appl. Math. Sci. **9** (2015), no. 31, 1517-1525.
19. A. V. Ustinov, *Korobov polynomials and umbral analysis,* (Russian) Chebyshevskiĭ Sb. **4** (2003), no. 4(8), 137-152.
20. N. L. Wang, H. Li, *Some identities on the Higher-order Daehee and Changhee Numbers,* Pure Appl. Math. J. **4** (2015), no. 5-1, 33-37.

1, Institute of Mathematics and Computer Science, Far Eastern Federal University, 690950 Vladivostok, Russia
E-mail address: dvdolgy@gmail.com

2, Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea
E-mail address: dskim@sogang.ac.kr

3, Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea
E-mail address: tkkim@kw.ac.kr

4, Department of Applied Mathematics, Pukyong National University, Busan 608-737, Republic of Korea
E-mail address: seo2011@pknu.ac.kr